

PATIENT NO: _____ DATE: _____ TEST OCCASION:

Baseline	Post ECT 3 or	Post ECT 4
----------	---------------	------------

BRIEF ECT COGNITIVE SCREEN (BECS)

The BECS is to be administered prior to commencement of ECT and at least 1 day after the 3rd or 4th ECT treatment.

- For each item below use a stopwatch to time the duration.
- Read the instructions for each item below.
- Calculate the total time and total errors for both items.
- Following post ECT 3 or post ECT 4 calculate the BECS Total Score out of 5 by adding together the scores for Total Errors and Total Time.

		Errors	Time (sec)
1.	<p><i>Now I would like for you to count backwards from twenty to one as quickly as you can. Like this, twenty, nineteen, eighteen, etc., all the way to one. Start now.</i></p> <p style="text-align: center;"> 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 </p>		
2.	<p><i>For this last one, I would now like you to say the months of the year backwards starting with December as quickly as you can. Start now.</i></p> <p style="text-align: center;"> December November October September August July June May April March February January </p>		
		Total Errors	Total Time (seconds)

CALCULATE DIFFERENCE

Post ECT 3	TOTAL ERRORS	TOTAL TIME	BECS TOTAL SCORE
<u>MINUS</u>			
Baseline			
DIFFERENCE			
SCORE			/5

Scoring

NB: Scoring only applies to positive values

If difference in errors < 1, score = 0
 If difference in errors = 1, score = + 1
 If difference in errors ≥ 2, score = + 2

If difference in time < 5, score = 0
 If difference in time ≥ 5, score = + 1
 If difference in time ≥ 8, score = + 2
 If difference in time ≥ 12, score = + 3

For more information regarding the BECS, see the following reference

Martin, D. M., Katalinic, N., Ingram, A., Schweitzer, I., Smith, D. J., Hadzi-Pavlovic, D., & Loo, C. K. (2013). A new early cognitive screening measure to detect cognitive side-effects of electroconvulsive therapy?. *Journal of psychiatric research*, 47(12), 1967-1974.